



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

May 13, 2016

Kristian Paul  
Regulatory Affairs Manager  
Bayer Crop Sciences, Environmental Science Division  
2 T.W. Alexander Drive  
RTP, NC 27709

Subject: Label Amendment – Adding container referral statement, deleting statement ‘Before using Armada 50 WDG on ornamental plants that are not listed on the label, test Armada 50 WDG on a small scale first’, adding ABN ‘Trigo’ and adding marketing claims.  
Product Name: Armada 50 WDG  
EPA Registration Number: 432-1513  
Application Date: 1-19-16  
Decision Number: 513365

Dear Kristian Paul:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. If you have any questions, please contact Nathan Mellor by phone at 703-347-8562, or via email at [mellor.nathan@epa.gov](mailto:mellor.nathan@epa.gov).

A handwritten signature in black ink, reading "Shaja B. Joyner". The signature is written in a cursive, flowing style.

Shaja B. Joyner, Product Manager 20  
Fungicide-Herbicide Branch  
Registration Division 7505P

# ARMADA™ 50 WDG

## [ABN: STRIKE PLUS™ 50 WDG]

---

**SECTION A: Turf uses, landscape and interiorscapes  
ornamental uses.**

**ACCEPTED**

**05/13/2016**

Under the Federal Insecticide, Fungicide  
and Rodenticide Act as amended, for the  
pesticide registered under  
EPA Reg. No. 432-1513

# ARMADA™ 50 WDG

## [ABN: STRIKE PLUS™ 50 WDG]

• Editorial Note – Marketing claim positioned here

For the enhancement of greener and more dense turfgrass and for the control of listed foliar, stem and root diseases of turfgrass including institutional, commercial and residential lawns, sod farms, sports fields, parks, municipal grounds and cemeteries, and for control of listed foliar and stem diseases of ornamentals and non-bearing fruit trees grown in interiorscapes (to include shopping malls, shopping centers, hotel lobbies) and residential and commercial landscapes.

Editorial Note – [Bracketed text] is optional

**ACTIVE INGREDIENT:**

Trifloxystrobin (CAS No. 141517-21-7).....	8.33%
Triadimefon (CAS No. 43121-43-3).....	41.67%

**OTHER INGREDIENTS:**..... 50.00%

**TOTAL:** 100.00%

EPA Reg. No. 432-1513

EPA Est. No. \_\_\_\_\_

**KEEP OUT OF REACH OF CHILDREN.**

## CAUTION

See [Back] [Side] Panel for First Aid Instructions and [Leaflet] [Booklet] for Complete Precautionary Statements and Directions for Use. (Note to reviewer: Location of additional precautionary statements, directions for use will vary between those listed, depending on container type/size.)

**For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577**

**For PRODUCT USE Information Call 1-800-331-2867**

### FIRST AID

<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything to an unconscious person.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth to-mouth if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-334-7577 for emergency medical treatment.	
<b>Note to Physician:</b> If ingested, induce emesis or lavage stomach. Treat symptomatically.	

# PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Causes moderate eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid breathing dust or spray mist. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

### Personal Protective Equipment (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves made of waterproof material such as neoprene, butyl rubber, barrier laminate or nitrile rubber when mixing/loading, when using handheld equipment or handheld nozzles.
- Shoes plus socks
- Chemical-resistant apron, when mixing/loading or cleaning spills or equipment.

See engineering controls for additional requirements.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Control Statements

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## User Safety Recommendations

Users should:

- Wash hands thoroughly with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinsate.

### Ground Water Advisory

The active ingredients in this product have properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow may result in ground water contamination.

Do not make applications when weather conditions favor drift from target area.

## PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

## DIRECTIONS FOR USE

**It is a violation of federal law to use this product in a manner inconsistent with its labeling.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

**Exception:** If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Long sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

## PRODUCT INFORMATION

### Mixing Procedures

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Agitation is necessary for proper dispersal of the product. Maintain agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

**ARMADA 50 WDG Alone:** Add 1/2 of the required water to the mix tank. With the agitator running, add the prescribed dose of **ARMADA 50 WDG** to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after **ARMADA 50 WDG** has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

**ARMADA 50 WDG+ Tank Mixtures:** Add 1/2 of the required amount of water to the tank mix. Start the agitation running before adding **ARMADA 50 WDG** and any tank mix partner(s). In general, tank mix partners must be added in this order: (1) products packaged in water soluble packaging, wettable powders, wettable granules (such as **ARMADA 50 WDG**), (2) liquid flowables, liquids; and (3) emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all of the mixture has been applied.

If using **ARMADA 50 WDG** in a tank mixture, observe all restrictions, directions for use, crop/sites, use rates, dilution ratios, precautions and limitations that appear on the tank mix product label. Do not exceed labeled dosage rate and follow the most restrictive label precautions and limitations. This product must not be mixed with any product that prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products and uses are registered.

**ARMADA 50 WDG** is compatible with most insecticide, fungicide and foliar nutrient products. However, the compatibility of **ARMADA 50 WDG** with tank mix partners must be tested before use.

To determine biological compatibility with other products, mix the products in the desired proportions, spray on target plants and observe for phytotoxicity seven days after the application.

**Use with additives:** Use of spray additives is not required. Any spray additive must be evaluated prior to use. Do not use in conjunction with organosilicate-based products or plant injury may occur. Label directions are based on data with no additives.

**Chemigation:** Do not apply this product through any type of irrigation system.

**Aerial Application:** Do not apply by aerial application.

**Resistance Management:** **ARMADA 50 WDG** contains the strobilurin class of chemistry, which exhibits no known cross-resistance to other chemical classes including sterol inhibitors, dicarboximides, benzimidazoles, anilinopyrimidines or phenylamides. However, certain fungal pathogens are known to develop resistance to products used repeatedly. Because resistance development cannot be predicted, the use of this product must conform to resistance management strategies. Such strategies may include rotating and/or tank mixing with products having different modes of action; or limiting the total number of applications per season. Bayer [OHP] encourages responsible product stewardship to ensure effective long-term control of the fungal diseases on this label. See specific directions in the turf section and the ornamentals section.

## TURFGRASS DISEASE CONTROL

**ARMADA 50 WDG** is a preventive and curative fungicide that may be applied to turf sites including institutional, commercial and residential lawns, sod farms, sports fields, parks, municipal grounds and cemeteries.

**ARMADA 50 WDG** is a broad spectrum fungicide for the control of dollar spot (*Sclerotinia homoeocarpa*), brown patch (*Rhizoctonia solani*), anthracnose (*Colletotrichum graminicola*), red thread (*Corticium fuciforme*), pink patch (*Limonomyces roseipellis*), leaf spot (*Bipolaris* spp., *Drechslera* spp.), gray leaf spot (*Pyricularia grisea*), rust (*Puccinia* spp.), pink snowmold (*Microdochium nivale*), microdochium patch (*Microdochium nivale*), southern blight (*Sclerotium rolfsii*), stripe smut (*Ustilago striiformis*), rapid blight and summer patch (*Magnaporthe poae*). When conditions are favorable for Pythium blight (*Pythium* spp.), mix a Pythium control fungicide with **ARMADA 50 WDG**.

1. **ARMADA 50 WDG** IS BEST USED IN A PREVENTIVE DISEASE CONTROL PROGRAM.
2. Apply listed rate in 1-2 gallons of water per 1,000 sq ft to ensure thorough coverage. For soil-borne diseases, such as summer patch and take-all patch, use sufficient water to reach the crown and upper root zone or irrigate immediately after application with sufficient water to move the active ingredient down into the crown and root zone of the turf.
3. Apply after mowing OR allow sprayed area to completely dry before mowing.
4. For control of both foliar and soil-borne diseases, allow sprayed area to completely dry before irrigation.
5. Under conditions optimum for high disease pressure, use the higher rate and the shorter interval.
6. For optimum turf quality and disease control, use **ARMADA 50 WDG** in conjunction with turf management practices that promote good health and optimum disease control.
7. Before use of any fungicide, proper diagnosis of the organism causing the disease is important. Use of diagnostic kits or other means of identification of the disease organism is essential to determine the best control measures.

### Plant Health

Apply **ARMADA 50 WDG** at 1.2 to 1.5 oz/1,000 sq ft to sod production of warm season grasses such as St. Augustine, centipede and Bermudagrass. The following plant health enhancements will be seen 30 to 60 days after application of **ARMADA 50 WDG**; visible leaf greenness and density. Sod laid under shade trees also benefits in turf quality when this application is made prior to transfer.

Apply **ARMADA 50 WDG** at 1.2-1.5 oz /1,000 sq ft 15-30 days pre-harvest for plant health benefits.

### Resistance Management for Turfgrass

Do not apply more than 2 sequential **ARMADA 50 WDG** fungicide applications for gray leaf spot control. Do not apply more than 3 sequential applications of **ARMADA 50 WDG** fungicide for all other diseases. Alternate with a fungicide having a different mode of action.

## Tank Mixes

For more broad-spectrum control **ARMADA 50 WDG** can be tank mixed with other fungicides. Check compatibility before tank mixing.

**ARMADA 50 WDG** CAN BE APPLIED TO ALL MAJOR TURFGRASS SPECIES. RATES ARE DEPENDENT UPON DISEASE AND THE ORGANISM THAT IT IS INCITED BY.

TURFGRASS DISEASES CONTROLLED WITH ARMADA 50 WDG			
DISEASE	ARMADA 50 WDG	OZ OF ARMADA 50 WDG PER 1,000 SQ FT	APPLICATION INTERVAL/TIMING
Dollar Spot	Apply when conditions are favorable for disease development.	0.6-1.5	14-28 days
Brown Patch	Apply when conditions are favorable for disease development.	0.6-1.5	14-28 days
Leaf Spot	Apply when conditions are favorable for disease development.	0.6	14 days
	For curative applications.	1.2-1.5	21-28 days
Anthracnose	Apply when conditions are favorable for disease development.	0.6-1.5	14-28 days
Summer Patch	Make initial application when the soil temperatures reach 65°F. Subsequent applications at 21-28 day interval. Do not make more than 3 sequential applications.	1.2-1.5	21-28 days
Gray Leaf Spot and Rapid Blight	Apply when conditions are favorable for disease development. For best results on Rapid Blight apply <b>ARMADA 50 WDG</b> at the highest rate.	0.6-1.5	14-28 days
Red Thread, Pink Patch	Apply when conditions are favorable for disease development.	0.6-1.5	14-28 days
Rust	Apply when conditions are favorable for disease development.	0.6-1.5	14-28 days
Southern Blight	Preventive applications only	0.6-1.5	Repeat at 14-day intervals.
Stripe Smut	Make the first application in the spring just before the turf breaks dormancy followed by a second application just prior to the summer heat stress period and a third application when the cool nighttime temperatures of the late summer or early fall return.	0.6	3 applications/season Spring, Early Summer and Early Fall
Microdochium Patch	Apply when conditions are favorable for disease development.	1.2-1.5	Fall-Early Spring
Pink Snow Mold	Apply one application in late fall before snow cover or early spring after snow melts. Do not apply on top of snow.	1.2-1.5	Late Fall
Take-All Patch	Apply 1 to 2 applications preventatively.	1.2-1.5	28 days
Necrotic Ring Spot	Apply 1 to 2 applications preventatively.	1.2-1.5	28 days
Large Patch	Apply 1 to 2 applications preventatively when conditions are favorable for disease development.	1.2-1.5	28 days
Fairy Ring	First application is prescribed in the spring when 5 day average daily soil temperatures taken at 2 inches are 60-65°F. The second application is prescribed 14 to 28 days later. Make applications in 2 gallons water/1,000 sq ft. Water in application with 1/8 to 1/4 inch of irrigation within 6-8 hours after application.  Tank mixing with Revolution wetting agent will decrease the fungicidal efficacy of <b>ARMADA 50 WDG</b> . Apply wetting agents 2 weeks after the <b>ARMADA 50 WDG</b> application on routine schedule for wetting agents.	1.2-1.5	14-28 days

For preventive applications, apply when conditions are favorable for disease. Use the higher rate/shorter interval (if a range is given) for curative applications or under severe disease conditions.

## Maximum Use Rates

For turfgrass, do not exceed a maximum of 207 oz of **ARMADA 50 WDG** per acre per year (4.76 oz/1000 sq ft).



## ORNAMENTAL DISEASE CONTROL

**ARMADA 50 WDG** is a broad-spectrum systemic fungicide for the control of listed foliar and stem diseases of ornamentals and non-bearing fruit trees grown in interiorscapes (to include shopping malls, shopping centers, hotel lobbies) and residential and commercial landscapes.

### Use restrictions for ARMADA 50 WDG on Ornamentals.

- Applications with hose-end sprayers are permitted only for outdoor use on ornamentals.
- Do not use **ARMADA 50 WDG** on bearing fruit trees. **ARMADA 50 WDG** may be used on non-bearing trees. Non-bearing trees are defined as trees that will not bear fruit until at least 1 year after treatment.

**Conversion Table for ARMADA 50 WDG: cup, gram, ounce**

1/4 Cup = 30.5 grams = 1.07 oz	1/3 Cup = 40.7 grams = 1.43 oz
1/2 Cup = 61 grams = 2.14 oz	3/4 Cup = 91.5 grams = 3.21 oz
1 Cup = 122 grams = 4.28 oz	

### Interiorscape, Commercial and Residential Landscape Applications

**Foliar Diseases:** **ARMADA 50 WDG** will control foliar diseases of ornamental plants when applied as a foliar spray to the plant species listed on this label. Apply 3 to 9 oz **ARMADA 50 WDG** per 100 gallons of spray solution. Apply as a full-coverage spray to the point of drip and repeat at 14 to 28 day intervals until the threat of disease is over. Begin applications when conditions are favorable for disease development and continue until the threat of disease is over. Use higher rates for shorter intervals under high disease pressure. Do not exceed a maximum of 207 oz of **ARMADA 50 WDG** per acre per year (4.76 oz/ 1000 sq ft).

PLANTS		
FLOWERING AND FOLIAGE PLANTS	ORNAMENTAL SHRUBS, TREES AND NON-BEARING FRUIT TREES	SHADE TREES
Ageratum	Amelanchier	Ash
Aster	Apple (non-bearing)	Aspen
Begonia	Azalea	Birch
Calendula	Barberry	Buckeye
Canna	Buckthorn	Chestnut
Carnation	Camellia	Cottonwood
Chrysanthemum	Cedar	Elm
Cineraria	Cherry(non-bearing)	Fir
Crassula	Crabapple	Locust
Dahlia	Crape myrtle	Maple
Daisy	Cypress, Leyland	Oak
Delphinium	Dogwood	Pine
Dianthus	Euonymus	Poplar
Four O'Clock	Gardenia	Russian Olive
Geranium	Hawthorn	Sycamore
Gerbera	Hemlock	Walnut
Grape Leaf Ivy	Holly	Willow
Hollyhock	Juniper	
Hydrangea	Leucothoe	
Iris	Lilac	
Kalanchoe	Mock-Orange	
Marigold	Mountain Laurel	
Nephthytis	Ninebark	
Pansy	Paulownia	
Petunia	Pear (non-bearing)	

Phlox	Photinia	
Poinsettia	Pittosporum	
Rose	Plum (non-bearing)	
Salvia	Potentilla	
Sedum	Privet	
Snapdragon	Pyracantha	
Sunflowers	Rhododendron	
Zinnia	Spirea	
	Viburnum	
	Vitex	

**NOTICE TO USER:** Plant tolerance to **ARMADA 50 WDG** has been found to be acceptable on all ornamentals that it has been tested on with the exception of Petunia, Violets and New Guinea impatiens. Due to the large number of species and varieties of ornamentals plants, it is impossible to test every one for tolerance to **ARMADA 50 WDG**. Neither the Manufacturer nor the Seller has determined whether or not **ARMADA 50 WDG** can be used safely on ornamental plants not specified on this label. The professional user must determine if **ARMADA 50 WDG** can be used safely prior to commercial use. In a small area test the labeled rates on a small number of plants for phytotoxicity prior to widespread use. Before using **ARMADA 50 WDG** in tank mixture with other products, test the mixture on a small number of plants for phytotoxicity prior to widespread use.

COMMON AND SCIENTIFIC NAMES OF DISEASES CONTROLLED BY ARMADA 50 WDG	
COMMON NAME	SCIENTIFIC NAME
Anthracnose	<i>Apiognomonia veneta</i> <i>Colletotrichum gloeosporioides</i> <i>Discula quercina</i> <i>Gloeosporium aridum</i> <i>Glomerella cingulata</i>
Black spot	<i>Diplocarpon rosae</i>
Downy mildew	<i>Peronospora</i> spp.
Leaf spot	<i>Cercospora</i> spp. <i>Entomosporium</i> spp. <i>Septoria</i> spp.
Powdery mildew	<i>Erysiphe</i> spp. <i>Microsphaera</i> spp. <i>Oidium</i> spp. <i>Podosphaera</i> spp. <i>Sphaerotheca</i> spp. <i>Uncinula</i> spp.
Rust	<i>Coleosporium</i> spp. <i>Gymnosporangium</i> spp. <i>Melampsoridium</i> spp. <i>Phragmidium</i> spp. <i>Puccinia</i> spp. <i>Uromyces</i> spp.
Scab	<i>Cladosporium</i> spp. <i>Venturia inaequalis</i>

## Maximum Use Rates in Ornamentals

For ornamental plants do not exceed a maximum of 207 oz of **ARMADA 50 WDG** per acre per year (4.76 oz/1000 sq ft).

## Conifer and Christmas Tree Applications

PRESCRIBED APPLICATIONS		
Christmas Trees (Except Concolor Fir)	Stem and Cone Rusts <i>Cronartium</i> spp. (Fusiform) <i>Peridermium</i> spp. <i>Endocronartium Harknessii</i> (Gall)  Tip blight <i>Sirococcus strobilinus</i>  Lophodermium Needlecast <i>Lophodermium pinestri</i>	9 oz/A
	<p>Apply specified dosage per acre or per 100 gal of water as a full coverage, dilute spray as needed. Full coverage of the trees is essential for maximum control. Use of nonionic spray adjuvant is prescribed. Time applications appropriately for the specific disease being controlled. A maximum of 207 oz of <b>ARMADA 50 WDG</b> may be applied per acre per season (4.76 oz/1000 sq ft).</p> <p>For rusts, begin applications when the needles break through the fascicle sheath. Make additional applications at 14 to 21-day intervals. Stop when galls become pale to white color.</p> <p>For tip blight, begin applications to coincide with bud break. Make two additional applications at 14-day intervals.</p> <p>For Lophodermium needlecast, begin applications to coincide with spore release, normally beginning in mid-July and ending in mid-October. Make applications at 21-day intervals. Extend interval to 28 days if spore release is light or dry weather is expected.</p>	
Pine (Seedlings) (Except California)	Pine Rust (Fusiform rust)	9 oz/A
	<p>Begin application prior to infection period and repeat as necessary at 14 to 21-day intervals depending upon disease pressure. Use lower rates in areas of low disease incidence and higher rates in areas of severe disease incidence. A maximum of 207 oz of <b>ARMADA 50 WDG</b> may be applied per acre per season (4.76 oz/1000 sq ft). A spreader-sticker is needed to help adhere spray solution to the pine trees.</p> <p>Do not apply <b>ARMADA 50 WDG</b> on recent grafted scions until one year after grafting.</p>	

## Restrictions for Applications on Conifers and Christmas Trees

1. For ground application, use a minimum of 50 gals of water/A.
2. To avoid spray drift, do not apply when conditions favor drift beyond the target area. Avoid spray overlap. For information on spray equipment and calibration, consult sprayer manufacturers and state recommendations. For specific local directions and spray schedules, consult the current state agricultural experiment station recommendations.
3. Use of spray additives are not required. Any spray additive must be evaluated prior to use. Do not use in conjunction with organosilicate-based products or plant injury may occur. Label directions are based on data with no additives.

## Resistance Management for Ornamentals

**ARMADA 50 WDG** contains a site-specific fungicide belonging to the strobilurin class of chemistry. Fungal pathogens are known to develop resistance to fungicides with a specific mode of action. When site-specific fungicides are used without a clear resistance management strategy, resistance development may be rapid, particularly with greenhouse use.

**ARMADA 50 WDG** exhibits cross-resistance to other strobilurins and fungicides within the **Strobilurin Type Action and Resistance** group (**STAR** compounds), but there is no known cross-resistance to fungicides of other classes including sterol inhibitors, dicarboximides, benzimidazoles, anilinoypyrimidines, phenylpyrroles or phenylamides.

Many plant pathogens have a history of fungicide resistance development. To minimize the risk of resistance development to **ARMADA 50 WDG**, the following practices are prescribed.

1. Use **ARMADA 50 WDG** preventively.
2. For Leaf Spots and diseases other than Powdery Mildew, Downy Mildew and Botrytis:

- Use no more than two (2) applications of **ARMADA 50 WDG** before rotating to another effective product that is not in the strobilurin class of chemistry for two (2) applications before rotating back to **ARMADA 50 WDG**.

**OR**

- Rotate to another fungicide of nonstrobilurin chemistry after each **ARMADA 50 WDG** application.

3. For Powdery Mildew, Downy Mildew and Botrytis:

- Between each **ARMADA 50 WDG** application, make two (2) applications of a fungicide of nonstrobilurin chemistry before rotating back to **ARMADA 50 WDG**.

**OR**

- Rotate to another fungicide of nonstrobilurin chemistry after each **ARMADA 50 WDG** application.

4. Make no more than four (4) foliar applications of **ARMADA 50 WDG** per season for each at risk pathogen.

## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Material that cannot be used as directed should be disposed of as directed below. In spill or leak incidents, keep unauthorized people away.

**Pesticide Disposal:** Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture or rinse water is a violation of federal law. If these wastes cannot be used according to label instruction, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

**Container Handling:** Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available or dispose of empty carton in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

## IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness plant injury or other property damage, or other unintended consequences may result because of factors beyond the control of Bayer CropScience LP as weather conditions, presence of other materials, or the manner of use or application.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE, HANDLING, OR APPLICATION OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

This product contains a chemical known to the state of California to cause developmental or reproductive harm.

### Net Contents:

\* ARMADA is a trademark of Bayer

Produced for








**Bayer Environmental Science**

**A Division of Bayer CropScience LP  
Bayer Environmental Science  
P.O. Box 12014, 2 T.W. Alexander Drive  
Research Triangle Park, North Carolina 27709**

[Optional Marketing Claims:]

Pictures:





# **ARMADA™ 50 WDG**

## **[ABN: STRIKE PLUS™ 50 WDG]**

---

**Section B: Nurseries, garden centers and greenhouse  
ornamental uses.**



# ARMADA™ 50 WDG

## [ABN: STRIKE PLUS™ 50 WDG, TRIGO]

---

• Editorial Note – Marketing claim positioned here

---

For the control of listed foliar and stem diseases of ornamentals including flowers, foliage plants, shrubs, shade trees and non-bearing fruit trees grown in nurseries, garden centers and greenhouses.

---

Editorial Note – [Bracketed text] is optional

---

**ACTIVE INGREDIENT:**

Trifloxystrobin (CAS No. 141517-21-7).....	8.33%
Triadimefon (CAS No. 43121-43-3).....	41.67%

<b>OTHER INGREDIENTS:</b> .....	50.00%
---------------------------------	--------

<b>TOTAL:</b>	100.00%
---------------	---------

EPA Reg. No. 432-1513

EPA Est. No. \_\_\_\_\_

---

**KEEP OUT OF REACH OF CHILDREN.**

### CAUTION

See [Back] [Side] Panel for First Aid Instructions and [Leaflet] [Booklet] for Complete Precautionary Statements and Directions for Use. (Note to reviewer: Location of additional precautionary statements, directions for use will vary between those listed, depending on container type/size.)

---

**For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577**

**For PRODUCT USE Information Call 1-800-331-2867**

---

### FIRST AID

<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>Hold eye open and rinse slowly and gently with water for 15 to 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If on skin or clothing:</b>	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15 to 20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice.</li> <li>Have person sip a glass of water if able to swallow.</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything to an unconscious person.</li> </ul>
<b>If inhaled:</b>	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-800-334-7577 for emergency medical treatment.	
<b>Note to Physician:</b> If ingested, induce emesis or lavage stomach. Treat symptomatically.	

# PRECAUTIONARY STATEMENTS

## HAZARDS TO HUMANS AND DOMESTIC ANIMALS

### CAUTION

Causes moderate eye irritation. Harmful if swallowed, inhaled or absorbed through the skin. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Avoid breathing dust or spray mist. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

### Personal Protective Equipment (PPE)

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves made of waterproof material such as neoprene, butyl rubber, barrier laminate or nitrile rubber when mixing/loading, when using handheld equipment or handheld nozzles.
- Shoes plus socks
- Chemical-resistant apron, when mixing/loading or cleaning spills or equipment.

See engineering controls for additional requirements.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Control Statements

When handlers use closed systems, enclosed cabs or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## User Safety Recommendations

Users should:

- Wash hands thoroughly with soap and water before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates. Do not apply directly to water or to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinsate.

### Ground Water Advisory

The active ingredients in this product have properties and characteristics associated with chemicals detected in ground water. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow may result in ground water contamination.

Do not make applications when weather conditions favor drift from target area.

## PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

## DIRECTIONS FOR USE

**It is a violation of federal law to use this product in a manner inconsistent with its labeling.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

## AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

**Exception:** If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water is:

- Long sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material
- Shoes plus socks

## NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses.

Do not enter treated areas without protective clothing until sprays have dried.

## PRODUCT INFORMATION

### Mixing Procedures

Prepare no more spray mixture than is needed for the immediate operation. Thoroughly clean spray equipment before using this product. Agitation is necessary for proper dispersal of the product. Maintain agitation throughout the spraying operation. Do not let the spray mixture stand overnight in the spray tank. Flush the spray equipment thoroughly following each use and apply the rinsate to a previously treated area.

**ARMADA 50 WDG Alone:** Add 1/2 of the required water to the mix tank. With the agitator running, add the prescribed dose of **ARMADA 50 WDG** to the tank. Continue agitation while adding the remainder of the water. Begin application of the solution after **ARMADA 50 WDG** has completely dispersed into the mix water. Maintain agitation until all of the mixture has been applied.

**ARMADA 50 WDG + Tank Mixtures:** Add 1/2 of the required amount of water to the tank mix. Start the agitation running before adding **ARMADA 50 WDG** and any tank mix partner(s). In general, tank mix partners must be added in this order: (1) products packaged in water soluble packaging, wettable powders, wettable granules (such as **ARMADA 50 WDG**), (2) liquid flowables, liquids; and (3) emulsifiable concentrates. Always allow each tank mix partner to become fully dispersed before adding the next product. Provide sufficient agitation while adding the remainder of the water. Maintain agitation until all of the mixture has been applied.

If using **ARMADA 50 WDG** in a tank mixture, observe all restrictions, directions for use, crop/sites, use rates, dilution ratios, precautions and limitations that appear on the tank mix product label. Do not exceed labeled dosage rate, and the most restrictive label precautions and limitations must be followed. This product must not be mixed with any product that prohibits such mixing. Tank mixtures or other applications of products referenced on this label are permitted only in those states in which the referenced products and uses are registered.

**ARMADA 50 WDG** is compatible with most insecticide, fungicide and foliar nutrient products. However, the compatibility of **ARMADA 50 WDG** with tank mix partners must be tested before use.

To determine biological compatibility with other products, mix the products in the desired proportions, spray on target plants and observe for phytotoxicity seven days after the application.

**Use with additives:** Use of spray additives is not required. Any spray additive must be evaluated prior to use. Do not use in conjunction with organosilicate-based products or plant injury may occur. Label directions are based on data with no additives.

**Chemigation:** Do not apply this product through any type of irrigation system.

**Aerial Application:** Do not apply by aerial application.

**Resistance Management:** **ARMADA 50 WDG** contains the strobilurin class of chemistry, which exhibits no known cross-resistance to other chemical classes including sterol inhibitors, dicarboximides, benzimidazoles, anilinopyrimidines or phenylamides. However, certain fungal pathogens are known to develop resistance to products used repeatedly. Because resistance development cannot be predicted, the use of this product must conform to resistance management strategies. Such strategies may include rotating and/or tank mixing with products having different modes of action; or limiting the total number of applications per season. Bayer [OHP] encourages responsible product stewardship to ensure effective long-term control of the fungal diseases on this label. See specific directions in the ornamentals section.

## ORNAMENTAL DISEASE CONTROL

**ARMADA 50 WDG** is a broad-spectrum systemic fungicide for the control of listed foliar and stem diseases of ornamentals including flowers, foliage plants, shrubs, shade trees and non-bearing fruit trees grown in nurseries, garden centers and greenhouses.

### Use restrictions for ARMADA 50 WDG on Ornamentals.

- Do not use **ARMADA 50 WDG** in residential greenhouses.
- Applications with hose-end sprayers are permitted only for outdoor use on ornamentals.
- Do not use **ARMADA 50 WDG** on bearing fruit trees. **ARMADA 50 WDG** may be used on non-bearing trees. Non-bearing trees are defined as trees that will not bear fruit until at least 1 year after treatment.

Conversion Table for ARMADA 50 WDG: cup, gram, ounce	
1/4 Cup = 30.5 grams = 1.07 oz	1/3 Cup = 40.7 grams = 1.43 oz
1/2 Cup = 61 grams = 2.14 oz	3/4 Cup = 91.5 grams = 3.21 oz
1 Cup = 122 grams = 4.28 oz	

### Garden Center and Nursery Applications

**Foliar Diseases:** **ARMADA 50 WDG** will control foliar diseases of ornamental plants when applied as a foliar spray to the plant species listed on this label. Apply 3 to 9 oz **ARMADA 50 WDG** per 100 gallons of spray solution. Apply as a full-coverage spray to the point of drip and repeat at 14 to 28 day intervals until the threat of disease is over. Begin applications when conditions are favorable for disease development and continue until the threat of disease is over. Use higher rates or shorter intervals under high disease pressure. Do not exceed a maximum of 207 oz of **ARMADA 50 WDG** per acre per year (4.76 oz/1000 sq ft).

Do not exceed a volume of 530 gallons of spray mix per acre per application at the highest application rate of 9 oz/100 gal.

PLANTS		
FLOWERING AND FOLIAGE PLANTS	ORNAMENTAL SHRUBS, TREES AND NON-BEARING FRUIT TREES	SHADE TREES
Ageratum	Amelanchier	Ash
Aster	Apple (non-bearing)	Aspen
Begonia	Azalea	Birch
Calendula	Barberry	Buckeye
Canna	Buckthorn	Chestnut
Carnation	Camellia	Cottonwood
Chrysanthemum	Cedar	Elm
Cineraria	Cherry(non-bearing)	Fir
Crassula	Crabapple	Locust
Dahlia	Crape myrtle	Maple
Daisy	Cypress, Leyland	Oak
Delphinium	Dogwood	Pine
Dianthus	Euonymus	Poplar
Four O'Clock	Gardenia	Russian Olive
Geranium	Hawthorn	Sycamore
Gerbera	Hemlock	Walnut
Grape Leaf Ivy	Holly	Willow
Hollyhock	Juniper	
Hydrangea	Leucothoe	
Iris	Lilac	
Kalanchoe	Mock-Orange	
Marigold	Mountain Laurel	
Nephthytis	Ninebark	
Pansy	Paulownia	

Petunia	Pear (non-bearing)	
Phlox	Photinia	
Poinsettia	Pittosporum	
Rose	Plum (non-bearing)	
Salvia	Potentilla	
Sedum	Privet	
Snapdragon	Pyracantha	
Sunflowers	Rhododendron	
Zinnia	Spirea	
	Viburnum	
	Vitex	

**NOTICE TO USER:** Plant tolerance to **ARMADA 50 WDG** has been found to be acceptable on all ornamentals that it has been tested on with the exception of Petunia, Violets and New Guinea impatiens. Due to the large number of species and varieties of ornamentals plants, it is impossible to test every one for tolerance to **ARMADA 50 WDG**. Neither the Manufacturer nor the Seller has determined whether or not **ARMADA 50 WDG** can be used safely on ornamental plants not specified on this label. The professional user must determine if **ARMADA 50 WDG** can be used safely prior to commercial use. In a small area test the labeled rates on a small number of plants for phytotoxicity prior to widespread use. Before using **ARMADA 50 WDG** in tank mixture with other products, test the mixture on a small number of plants for phytotoxicity prior to widespread use.

COMMON AND SCIENTIFIC NAMES OF DISEASES CONTROLLED BY ARMADA 50 WDG	
COMMON NAME	SCIENTIFIC NAME
Anthracnose	<i>Apiognomonia veneta</i> <i>Colletotrichum gloeosporioides</i> <i>Discula quercina</i> <i>Gloeosporium aridum</i> <i>Glomerella cingulata</i>
Black spot	<i>Diplocarpon rosae</i>
Downy mildew	<i>Peronospora</i> spp.
Leaf spot	<i>Cercospora</i> spp. <i>Entomosporium</i> spp. <i>Septoria</i> spp.
Powdery mildew	<i>Erysiphe</i> spp. <i>Microsphaera</i> spp. <i>Oidium</i> spp. <i>Podosphaera</i> spp. <i>Sphaerotheca</i> spp. <i>Uncinula</i> spp.
Rust	<i>Coleosporium</i> spp. <i>Gymnosporangium</i> spp. <i>Melampsoridium</i> spp. <i>Phragmidium</i> spp. <i>Puccinia</i> spp. <i>Uromyces</i> spp.
Scab	<i>Cladosporium</i> spp. <i>Venturia inaequalis</i>

## Greenhouse Applications

Except as noted for specific diseases, mix 1.2 to 2.4 ounces of **ARMADA 50 WDG** in 100 gallons of water and apply as a full coverage foliage spray to the point of drip. Do not exceed a volume of 530 gallons of spray mix per acre at the highest dose rate of 9 oz/100 gal.

Winter Use –1.2 oz of **ARMADA 50WDG**.

Summer Use – 2.4 oz of **ARMADA 50 WDG**.

Mix specified amount of **ARMADA 50 WDG** in 100 gallons of water and apply in a spray application to the point of drip. Intervals between applications must be no shorter than 30 days to avoid flower stalk length reduction. Excessive rates or applications may result in a shortening of the flower stalk.

PLANT	DISEASE		APPLICATION RATE	APPLICATION TIMING	INTERVAL BETWEEN APPLICATIONS
	COMMON NAME	SCIENTIFIC NAME			
African violet Cineraria Crassula Gerbera Grape leaf Ivy Hydrangea Kalanchoe Poinsettia	Powdery Mildew	<i>Erysiphe</i> spp., <i>Microsphaera</i> spp., <i>Oidium</i> spp., <i>Phyllactinia</i> spp. <i>Podosphaera</i> spp., <i>Sphaerotheca</i> spp. <i>Uncinula</i> spp.	1.2-2.4 oz /100 gal	Apply when conditions are favorable for disease development	No shorter than 30 days to avoid flower stalk length reduction*
Azalea	Powdery Mildew	<i>Erysiphe</i> spp., <i>Microsphaera</i> spp., <i>Oidium</i> spp., <i>Phyllactinia</i> spp. <i>Podosphaera</i> spp., <i>Sphaerotheca</i> spp. <i>Uncinula</i> spp.	1.2-2.4 oz /100 gal	Apply when conditions are favorable for disease development	No shorter than 30 days to avoid flower stalk length reduction*
	Anthracnose /Flower blight	<i>Ovulinia</i> spp	4.8-9 oz / 100 gal	Begin applications at the expanded bud stage (color showing)	7 to 14-day intervals as needed dependent upon bloom periods.
	Leaf spot/Blights	<i>Exobasidium</i> spp.	2.4 oz / 100 gal	Begin applications at bud break	10 days as needed
Calendula Carnation Chrysanthemum Daisy Geranium Snapdragon	Powdery Mildew	<i>Erysiphe</i> spp., <i>Microsphaera</i> spp., <i>Oidium</i> spp., <i>Phyllactinia</i> spp. <i>Podosphaera</i> spp., <i>Sphaerotheca</i> spp. <i>Uncinula</i> spp.	1.2-2.4 oz /100 gal	Apply when conditions are favorable for disease development	No shorter than 30 days to avoid flower stalk length reduction*
	Rust	<i>Gymnosporangium</i> spp.	4.8-9 oz / 100 gal	Begin applications at the expanded bud stage (color showing)	7-14 days as needed dependent upon bloom period
		<i>Melampsoridium</i> spp.			
		<i>Phragmidium andersonii</i>			
		<i>Uromyces</i> spp.			
		<i>Melampsora farlowii</i> <i>Uredinopsis mirabilis</i> spp.	9 oz / 50 gal	Apply in spring to shoots in the upper whorl of susceptible pine species	Make a single application per year
		<i>Melampsora pinitorqua</i>	9 oz (plus sufficient spreader sticker) / 100 gal	Apply when conditions are favorable for disease development	As needed basis during the early part of the season*
		<i>Cronartium</i> spp. <i>Peridermium</i> spp.			

		<i>Coleosporium spp.</i> <i>Puccinia spp.</i>	4.8-9 oz / 100 gal	Apply in early spring as growth starts	14-21 days until new growth is fully expanded
Rose	Powdery Mildew	<i>Erysiphe spp.</i> , <i>Microsphaera spp.</i> , <i>Oidium spp.</i> , <i>Phyllactinia spp.</i> , <i>Podosphaera spp.</i> , <i>Sphaerotheca spp.</i> , <i>Uncinula spp.</i>	1.2-2.4 oz /100 gal	Apply when conditions are favorable for disease development	No shorter than 30 days to avoid flower stalk length reduction*
	Black spot	<i>Diclocarpon rosae</i>	1.2 (winter)- 2.4(summer) / 100 gal	Apply when conditions are favorable for disease development	No shorter than 30 days to avoid flower stalk length reduction*

\* Excessive rates or applications may result in a shortening of the flower stalk.

### Maximum Use Rates in Ornamentals

For ornamental plants do not exceed a maximum of 207 oz of **ARMADA 50 WDG** per acre per year (4.76 oz/1000 sq ft). Do not exceed a volume of 530 gallons of spray mix per acre per application at the highest application rate of 9 oz/100 gal.

### Resistance Management for Ornamentals

**ARMADA 50 WDG** contains a site-specific fungicide belonging to the strobilurin class of chemistry. Fungal pathogens are known to develop resistance to fungicides with a specific mode of action. When site-specific fungicides are used without a clear resistance management strategy, resistance development may be rapid, particularly with greenhouse use.

**ARMADA 50 WDG** exhibits cross-resistance to other strobilurins and fungicides within the **Strobilurin Type Action and Resistance** group (**STAR** compounds), but there is no known cross-resistance to fungicides of other classes including sterol inhibitors, dicarboximides, benzimidazoles, anilinopyrimidines, phenylpyrroles or phenylamides.

Many plant pathogens have a history of fungicide resistance development. To minimize the risk of resistance development to **ARMADA 50 WDG**, the following practices are prescribed.

1. Use **ARMADA 50 WDG** preventively.
2. For Leaf Spots and diseases other than Powdery Mildew, Downy Mildew and Botrytis:
  - Use no more than two (2) applications of **ARMADA 50 WDG** before rotating to another effective product that is not in the strobilurin class of chemistry for two (2) applications before rotating back to **ARMADA 50 WDG**.

**OR**

  - Rotate to another fungicide of nonstrobilurin chemistry after each **ARMADA 50 WDG** application.
3. For Powdery Mildew, Downy Mildew and Botrytis:
  - Between each **ARMADA 50 WDG** application, make two (2) applications of a fungicide of nonstrobilurin chemistry before rotating back to **ARMADA 50 WDG**.

**OR**

  - Rotate to another fungicide of nonstrobilurin chemistry after each **ARMADA 50 WDG** application.
4. Make no more than four (4) foliar applications of **ARMADA 50 WDG** per season for each at risk pathogen



## STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

Handle and open container in a manner as to prevent spillage. If the container is leaking or material spilled for any reason or cause, carefully sweep material into a pile. Refer to Precautionary Statements on label for hazards associated with the handling of this material. Do not walk through spilled material. Material that cannot be used as directed should be disposed of as directed below. In spill or leak incidents, keep unauthorized people away.

**Pesticide Disposal:** Pesticide wastes may be toxic. Improper disposal of unused pesticide, spray mixture or rinse water is a violation of federal law. If these wastes cannot be used according to label instruction, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance in proper disposal methods.

**Container Handling:** Non-refillable container. Do not reuse or refill this container. Triple rinse container (or equivalent) promptly after emptying. Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container 1/4 full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Offer for recycling, if available or dispose of empty carton in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

## IMPORTANT: READ BEFORE USE

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness plant injury or other property damage, or other unintended consequences may result because of factors beyond the control of Bayer CropScience LP as weather conditions, presence of other materials, or the manner of use or application.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR CONSEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES RESULTING FROM THE USE, HANDLING, OR APPLICATION OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

This product contains a chemical known to the state of California to cause developmental or reproductive harm.

### Net Contents:

\* ARMADA, is a trademark of Bayer

\* STRIKE PLUS 50 WDG is a trademark of OHP

Produced for



Bayer Environmental Science

A Division of Bayer CropScience LP

Bayer Environmental Science

P.O. Box 12014, 2 T.W. Alexander Drive

Research Triangle Park, North Carolina 27709

ARMADA 50 WDG (PENDING) 01/12/2016, 01/15/2016, 01/19/2016, 05/06/2016



[Optional Marketing Claims:]

- Fungicide
- Ornamental fungicide
- For nursery and greenhouse use

Pictures:

